

## PERIPHERAL STIMULATION FOR TREATMENT OF TRIGEMINAL POSTHERPETIC NEURALGIA AND TRIGEMINAL POSTTRAUMATIC NEUROPATHIC PAIN: A PILOT STUDY

Mark D. Johnson, M.D.,  
Ph.D.

Department of Neurological  
Surgery, Harvard Medical School  
and Brigham and Women's  
Hospital, Boston, Massachusetts

Kim J. Burchiel, M.D.

Department of Neurological  
Surgery, Oregon Health & Science  
University, Portland, Oregon

**Reprint requests:**

Kim J. Burchiel, M.D.,  
Department of Neurological  
Surgery,  
Oregon Health & Science  
University, Mail Code 4572,  
3131 SW Sam Jackson Park Road,  
Portland, OR 97239.  
Email: kburch@ohsu.edu

Received, August 5, 2004.

Accepted, March 4, 2004.

**OBJECTIVE:** Trigeminal neuropathic pain (TNP) after facial trauma or herpes zoster infection is often refractory to treatment. Peripheral nerve stimulation has been used to treat occipital neuralgia, however, efficacy in controlling facial TNP or postherpetic neuralgia is unknown. A retrospective case series of patients who underwent subcutaneous placement of stimulating electrodes for treatment of V<sub>1</sub> or V<sub>2</sub> TNP secondary to herpetic infection or facial trauma is presented.

**METHODS:** Ten patients received implanted subcutaneous pulse generators and quadripolar electrodes for peripheral stimulation of the trigeminal nerve supraorbital or infraorbital branches. Long-term treatment results were determined by retrospective review of medical records (1998-2003) and by independent observers interviewing patients using a standard questionnaire. Surgical complication rate, preoperative symptom duration, degree of pain relief, preoperative and postoperative work status, postoperative changes in medication usage, and overall degree of therapy satisfaction were assessed. Mean follow-up was 26.6 ± 4.7 months.

**RESULTS:** Peripheral nerve stimulation provided at least 50% pain relief in 70% of patients with TNP or postherpetic neuralgia. Medication use declined in 70% of patients, and 80% indicated that they were mostly or completely satisfied with treatment overall. There were no treatment failures (<50% pain relief and a lack of decrease in medication use) in the posttraumatic group, and two failures (50%) occurred in the postherpetic group. The complication rate requiring reoperation was 30%.

**CONCLUSION:** Peripheral nerve stimulation of the supraorbital or infraorbital branches of the trigeminal nerve is an effective method for relief of TNP after facial trauma or herpetic infection. A prospective trial using this novel approach to treat these disorders is thus warranted.

**KEY WORDS:** Peripheral nerve stimulation, Postherpetic neuralgia, Posttraumatic neuropathic pain, Trigeminal

Neurosurgery, 55:135-142, 2004

DOI: 10.1227/00006123-200407000-00010

www.neurosurgery-online.com

Unlike classic trigeminal neuralgia, facial pain occurring after traumatic injury or herpes zoster infection of the facial branches of the trigeminal nerve is a medical problem that is often very difficult to treat (3). Tricyclic antidepressants, topical anesthetics or capsaicin, intrathecally administered corticosteroids or local anesthetics, and gabapentin have all been shown to have limited efficacy in the treatment of postherpetic neuralgia (1, 2, 5, 10). Microvascular decompression is generally ineffective in this setting. Peripheral neurectomy and percutaneous gangliolysis have also been attempted in certain patients, but again with limited success

(1, 2). Motor cortex stimulation and stereotactic trigeminal neurectomy have been tried with reported success in 50 to 70% of patients (6, 9, 11-13, 15, 16). Patients are often quite disabled by their symptoms; consequently, a safer, more accessible, and more effective treatment is needed.

Peripheral nerve stimulation has been used to treat occipital neuralgia, although detailed series describing the efficacy and long-term outcome of this approach are lacking. Moreover, there are no reports in the medical literature describing the use or long-term efficacy of subcutaneous stimulation of branches of the trigeminal nerve for the treatment of facial pain occur-

ing after traumatic injury or herpes zoster infection. To determine whether peripheral trigeminal nerve stimulation might be an effective therapy for these disorders, either subcutaneous or intraneural stimulating electrodes were implanted in a cohort of patients who presented with refractory neuropathic facial pain (first in the territory of the first or second branch of the trigeminal nerve) secondary to traumatic injury or herpes infection. Medical records were reviewed retrospectively, and patient outcomes were re-evaluated to assess treatment outcome.

## PATIENTS AND METHODS


### Surgical Technique

Informed consent for surgery was obtained before all surgical procedures. Patients were taken to the operating room and placed in a supine position. The face and neck were prepped and draped in a sterile manner. In most patients, a general anesthetic was used. In a minority of patients, however, the procedure was performed under a local anesthetic with the patient awake. With the head turned to expose the temporal and cervical region of the affected side, a preauricular vertical incision was made. Under fluoroscopic guidance, a four contact Triax Quad stimulating electrode (Model 3422A, Medtronic, Inc., Minneapolis, MN) was advanced concurrently into the region of either the supraorbital or infraorbital foramen with a Tuohy needle (Fig. 1). This electrode contained four linearly arranged 2 mm electrode contacts that were spaced 6 mm apart. The proximal end of the electrode was then connected to an extension cable, which was subsequently tunneled subcutaneously to an exit point in the upper cervical region. A trial period of 2 to 7 days (mean 4.5 days), during which the patient was given a temporary external pulse generator and was taught to self-adjust the stimulation parameters to maximize pain relief. Hospital staff assisted the patients in selecting parameters that provided direct coverage of their painful areas. Patients undergoing a 7-day trial were kept in hospital throughout the trial period. Those patients who underwent a longer trial period were discharged to home and brought back on an outpatient basis for implantation of the pulse generator.

Patients who experienced significant therapeutic benefit were taken back to the operating room for implantation of an Impl. Pulse Generator (Model 3425; Medtronic, Inc.). Patients were again placed in the supine position and analgesized with general anesthesia. The temporary extension cable was removed, and a permanent extension cable was tunneled subcutaneously from an infraorbital or supraorbital foramen to the supraorbital or infraorbital site, where it was connected to the proximal end of the stimulating extension cable tunneled to the preauricular site, where it was connected to the extension cord (Fig. 2). The subcutaneous tunnel was tunneled to the other end of the extension cable and placed into a surgically prepared infraorbital or supraorbital pocket above the preauricular incision. Postoperative visits were scheduled within 14 days of the second surgical procedure.



**FIGURE 1.** Approach of patient to supraorbital and infraorbital foramina. Image A shows and image B shows the approach of supraorbital (A) and infraorbital (B) quadrupole stimulating electrodes for the treatment of neuropathic pain and posttraumatic neuroma pain. The lateral view (B) also demonstrates the retroauricular location of the proximal and extension cable which was tunneled subcutaneously to an infraorbital or supraorbital foramen in the patient (top).



### Data Collection and Analysis

Medical records from 1995 to 2005 were reviewed to identify patients who had undergone placement of subcutaneous stimulating electrodes for the treatment of facial pain occurring after traumatic injury or herpes infection. A trial case form was then developed to assess preoperative symptom duration, patterns of stimulator use, postoperative changes in medication usage, degree of pain relief, cooperation and postoperative work status, and overall level of satisfaction with the therapy. Independent observers who were not members of the surgical team conducted telephone interviews to obtain follow-up data. Information on surgical complications was obtained from medical records and from patient interviews. Statistical analysis was performed with StatView (SAS Institute, Inc., Cary, NC), a commercially available software package. Data are presented as mean  $\pm$  standard error of the mean. The procedures and data analysis for this study were performed in accordance with concurrent Oregon Health & Science University Institutional Review Board research guidelines.

## RESULTS

### Patient Demographics

A total of 77 patients who had undergone surgical placement of a subcutaneous stimulating electrode in the facial region

were identified. With one exception, all patients presented with neurovascular facial pain in the context of a history of facial trauma or herpes zoster infection affecting the face. The exception was a patient with spontaneous idiopathic facial pain that was refractory to numerous forms of medical and surgical therapy. Typical symptoms included allodynia, hyperesthesia or hypersensitivity, and chronic burning sensations in the distribution of the supraorbital or infraorbital nerves. No patient had comorbid anesthesia in the involved dermatome. All of the patients had undergone unsuccessful trials of one or more other forms of therapy before surgery, including antidepressant and tricyclic antidepressant (e.g., gabapentin or amitriptyline), topical anesthetics, peripheral neurectomy, percutaneous trigeminal gangliolysis, or microvascular decompression.

Eight of the patients were affected in the distribution of the supraorbital nerve, and three were affected in the distribution of the infraorbital nerve. Only patients with pain in the  $V_1$  or  $V_2$  distribution were considered for implantation of a nerve stimulator. The authors elected not to place stimulating electrodes in the  $V_3$  territory because of concerns about lead breakage caused by repeated movement of the mandible. One of the 11 patients who had sustained a traumatic infraorbital nerve injury failed to obtain significant relief after a 1-week trial period of peripheral nerve stimulation, consequently, no pulse generator was implanted.

Of the remaining 10 patients who received a permanent generator implant, 7 were men and 3 were women, yielding a male-to-female ratio of 2.3:1. The mean age at the time of implantation was  $52.2 \pm 6.2$  years (range, 35–66 yr). As shown in Figure 1, 40% of the patients experienced postoperative neuropathic pain, 40% postherpetic neuralgia, and 20% atypical facial pain

that was unresponsive to traditional treatment modalities for trigeminal neuralgia (including anticonvulsant therapy, gamma nucleus gangliolysis, and microvascular decompression). Eight (80%) of the 10 patients were contacted for completion of a standardized questionnaire. Data for the remaining 2 patients (with posttraumatic neuropathic pain and 1 with postherpetic neuralgia) were gleaned from medical records of postoperative clinic visits. The mean duration of symptoms before surgery was  $47.5 \pm 13.6$  months (range, 7–111 mo). Data characterizing the patient population are summarized in Table 1 and Figure 2.

### Therapeutic Efficacy

Several parameters were assessed to determine whether peripheral trigeminal nerve stimulation was effective in relieving postoperative or preoperative neuropathic pain. These included degree of pain relief, preoperative and postoperative work status, changes in medication usage, and overall satisfaction rating.

Assessment of the degree of pain relief afforded by peripheral stimulation of the supraorbital or infraorbital branches of the trigeminal nerve revealed that 70% of patients experienced a 50% or greater degree of pain relief after undergoing this procedure. Overall, the mean degree of pain relief was  $58 \pm 10\%$  (range, 0–100%), with a median of approximately 75%. The distribution of the results is shown in Figure 3. There was no correlation between the degree of preoperative pain relief and patient age, sex, or duration of preoperative symptoms.

Seven of the 10 patients reported a decrease in medication use after surgery, an effect that was statistically significant ( $P < 0.02$ , Wilcoxon  $T$ -test). Two patients experienced no change in their medication use after surgery, and only one patient reported an increase in medication use (Fig. 4). This patient (Patient 4) had toxic pallor of mucous membranes, which may explain why she experienced less than 50% pain relief and an increase in medication use despite peripheral trigeminal stimulation. As expected, there was a close correlation between changes in medication use and the degree of pain relief experienced after surgery. Without exception, patients with at least a greater pain relief experienced an overall decrease in medication use, whereas patients with less than 50% pain relief had either no change or an increase in their medication use.

Preoperative and postoperative work status was also determined in an effort to assay the efficacy of peripheral trigeminal nerve stimulation. Before surgery, 60% of the patients were employed, and 40% were unemployed. Half of those who were unemployed, however, were in the eighth decade of life and well beyond retirement age. Thus, six of the eight patients who were under retirement age were employed before surgery. After implantation of the trigeminal nerve stimulating electrodes, seven of the eight patients returned to work. This small increase in the number of employed patients after surgery indicates that the surgical procedure had no deleterious effect on work status.

Degree of patient satisfaction was used as another measure of therapeutic efficacy. As shown in Figure 2, 90% of patients indicated that they were mostly or completely satisfied with





FIGURE 1. A, pie diagram illustrating the distribution of pain types for facial nerve among the patients in this study and preoperative burden of facial stimulating electrodes and generator. B, graph illustrating the distribution of preoperative symptoms relative to the duration of peripheral nerve stimulation with a permanent generator. The mean duration of preoperative symptoms was  $47.5 \pm 13.6$  months.



**FIGURE 3.** A, graph illustrating degree of pain relief achieved by peripheral trigeminal stimulation as reported by patients at the time of last follow-up. Seven of 10 patients reported a 50% or greater degree of pain relief after surgery. B, graph illustrating the percentage change in medication usage after implantation of trigeminal nerve stimulators reported by patients at the time of last follow-up. The majority of patients reported a decrease in medication use after surgery. C, graph showing preoperative and postoperative work status. Work status was necessary to determine the effect of trigeminal stimulation on functional outcomes. Two of the 10 patients in the study were beyond retirement age before surgery. Thus, only 8 patients of working age returned to work after surgery. D, graph showing the degree of patient satisfaction. Patient ratings of satisfaction were used to evaluate degree of patient pain relief and satisfaction of trigeminal nerve stimulator treatment at the time of last follow-up. 80% of patients indicated that they were totally or completely satisfied with the procedure.

the procedure. In the majority of patients, the degree of satisfaction correlated closely with the degree of pain relief and changes in medication use. In one case, however, a patient who reported only 20% pain relief and an increase in medication usage nevertheless reported that she was "extremely satisfied" with the procedure. This finding emphasizes the importance of obtaining both singular and global measures of efficacy when attempting to determine the usefulness of pain relief measures.

#### Durability of Peripheral Trigeminal Nerve Stimulation

Studies in other systems have indicated that the efficacy of peripheral nerve stimulation can decrease over time.<sup>11</sup> However, the durability of the therapeutic effect over time was determined. Patients were first questioned regarding their patterns of use of the generators. All of the patients used their stimulators at least 50% of the time, with 6 of the 10 patients keeping their turned on 100% of the time.

A necessary criterion for permanent implantation of the stimulating electrode and generator was that patients receive at least

50% pain relief from the device during the initial 1-year period. Using a level of 50% pain relief or greater as a marker of therapeutic efficacy, the number of patients deriving therapeutic benefit from the device as a function of time after implantation of the pulse generator was determined and plotted. As shown in Figure 6, the number of patients reporting at least 50% pain relief gradually decreased with longer periods of follow-up. However, 80% of patients continued to experience at least 50% pain relief after 24 months of follow-up.

#### Postherpetic versus Posttraumatic Neuropathic Pain

An examination of whether there were differences in therapeutic efficacy between trigeminal nerve stimulator for postherpetic or posttraumatic neuropathic pain was conducted. All the patients who had pain secondary to traumatic injury experi-

enced 50% or greater pain relief, compared with only half of these patients with postherpetic pain as the cause of their pain (Fig 3). There were similar differences between postherpetic and postherpetic trigeminal pain patients in terms of the effects of surgery on medication use and on overall satisfaction rating. Of the patients with posttraumatic neuropathic facial pain ( $n = 5$ ), 100% experienced a decrease in medication use after surgery, compared with 50% ( $n = 4$ ) of those with postherpetic pain. Similarly, all of the patients with posttraumatic facial pain reported that they were mostly or completely satisfied with the procedure, compared with only half of those with postherpetic pain. Although these differences were not statistically significant ( $P < 0.05$ , Student's *t* test), the findings suggest that a difference may exist between the ability of peripheral trigeminal nerve stimulation to treat posttraumatic and postherpetic neuropathic pain. However, the fact that half of the patients with postherpetic neuropathic pain experienced greater than 50% pain relief and a decrease in medication use after surgery indicates that this procedure can be an effective treatment in a subpopulation of patients with this difficult disorder.

TABLE 1. Characteristics of patient population\*

| Patient no. | Age/sex (yr) | Diagnosis                                  | Preoperative treatment | Procedure               | Complications         |
|-------------|--------------|--|------------------------|-------------------------|-----------------------|
| 1           | 60/F         | Traumatic right supraorbital neuralgia     | §, ¶, §                | Supraorbital stimulator | Wound breakdown       |
| 2           | 60/M         | Post-herpetic right supraorbital neuralgia | §, ¶                   | Supraorbital stimulator | None                  |
| 3           | 60/F         | Post-herpetic right supraorbital neuralgia | §, ¶, §, §             | Supraorbital stimulator | None                  |
| 4           | 60/F         | Post-herpetic left supraorbital neuralgia  | §, ¶, §, §, §, §       | Supraorbital stimulator | Wound breakdown       |
| 5           | 60/M         | Post-herpetic left supraorbital neuralgia  | §, ¶, §, §             | Supraorbital stimulator | Short extension cable |
| 6           | 60/F         | Traumatic right supraorbital neuralgia     | §, ¶, §, §             | Supraorbital stimulator | None                  |
| 7           | 60/M         | Post-herpetic left supraorbital neuralgia  | §, ¶, §                | Supraorbital stimulator | None                  |
| 8           | 60/F         | Traumatic left supraorbital neuralgia      | §, ¶, §, §             | Supraorbital stimulator | None                  |
| 9           | 60/F         | Traumatic left infraorbital neuralgia      | §, ¶, §, §, §          | Infraorbital stimulator | None                  |
| 10          | 60/F         | Traumatic right infraorbital neuralgia     | §, ¶, §                | Infraorbital stimulator | None                  |

\*The patient population is similar to an interventional study conducted by the author in the United States. Current practice includes the use of a percutaneous approach to implantation of the stimulator with accompanying physical therapy, behavioral therapy, and pharmacologic therapy. Patients undergoing physical therapy were not included.

### Complications

There were three complications requiring surgical intervention for an overall complication rate of 30%. All were associated with the retroauricular position of the connector and extension lead. In two patients, wound breakdown developed over the connector. Both required surgical revision. In one of these two patients, soft tissue infection led to the explant and subsequently reimplanted at a later date. The third patient complained of discomfort associated with tension on the extension lead when turning his head. It was taken back to the operating suite electively, where the extension lead was lengthened.




FIGURE 4. Graph illustrating the durability of peripheral trigeminal nerve stimulation, expressed as the number of patients reporting at least 50% pain relief as a function of time after surgery. The slope of the curve demonstrates a gradual decrease over time in the number of patients with good pain relief. By extrapolation, the number of months 100% of patients would experience 100% pain relief or greater would occur at approximately 60 months.

### DISCUSSION

Facial neuroathic pain of peripheral origin characteristically has several important features, including a precipitating pathogenesis such as disease (e.g., herpes zoster infection) or trauma, a delay period of days to months before onset, and typical symptoms such as burning paroxysmal or constant pain and dyesthesia that often occur in an area of complete sensory deficit (3,7). These features can be distinguishing if noted by the more commonly encountered trigeminal neuralgia, which is usually spontaneous in origin, is characterized by sharp lancinating pains, and is often the result of intracranial vascular compression of the trigeminal root. Facial neuropathic pain of peripheral origin should be distinguished




FIGURE 5. Bar graph depicting the efficacy of peripheral trigeminal stimulation for relief of pain secondary to either peripheral or neuroathic pain or peripheral neuropathy. There was a trend toward greater efficacy for the treatment of neuroathic neuroathic pain, although this difference was not statistically significant ( $P < 0.05$ ).

from anesthesia dolorosa, which is a deafferentation type of pain that is characterized by marked sensory loss in the periphery and that is probably caused by activation of abnormal central pain mechanisms (8).

The pathophysiology of facial neuropathic pain of trigeminal origin is poorly understood, although several studies suggest that peripheral mechanisms may not be the cause and that these may differ from patient to patient and may depend on the underlying cause of the pain (5, 7, 14). Increased tactile and temperature thresholds, as well as abnormal summation of pain, have been observed in patients with traumatic trigeminal nerve injury but not in patients with trigeminal pain of spontaneous origin (9). Postherpetic neuralgia is most likely has a different underlying pathophysiology that involves abnormal ties of both peripheral and central origin (14). Moreover, studies indicate that facial postherpetic neuralgia and truncal postherpetic neuralgia may be two separate pathophysiological entities (14).

As discussed previously, the treatment of facial neuropathic pain occurring after herpetic infection or traumatic injury is difficult. Evidence suggests that the early use of antiviral agents may aid in preventing the occurrence of postherpetic neuralgia (10). Once the disorder develops, however, it is often refractory to therapy. In this study, evidence for a novel approach to treatment of this disorder is presented. The data shown here indicate that the use of subcutaneous stimulating electrodes for peripheral stimulation of the supraorbital or infraorbital nerves yielded good to excellent pain control in 70% of patients, as measured by at least 30% pain relief, a decrease in medication usage, and high overall satisfaction ratings. The therapeutic benefits were durable, with 70% of patients reporting good to excellent results up to 4 years after the initiation of therapy.

One advantage of the procedure described here is that it uses hardware that is currently available for clinical use. The Paces-Quad stimulating electrode and the 5 Pulse Generator system are commonly used for epidural spinal cord stimulation. The device has been adapted for use in the facial region with good results. In fact, the results reported here closely parallel those for use of the Paces-Quad stimulating electrode in spinal cord stimulation, which produced 50% pain relief in 77% of patients after 1 year of therapy (4). Unlike spinal cord stimulation, which had no effect on medication usage, peripheral trigeminal nerve stimulation resulted in a decrease in medication usage in 71% of patients. The results of this study may also be compared with those reported for direct stimulation of the gasserian ganglion (17) for the treatment of postherpetic neuralgia, facial pain (70% good to excellent outcomes) and peripheral nerve stimulation (17) for the treatment of postherpetic pain (75% with good to excellent outcomes).

The complication rate of 30% observed in the present study is somewhat higher than the 12% reported for use of stimulating electrodes for spinal cord stimulation (4) but is comparable to that of 30 to 40% reported for stimulation of the gasserian ganglion (17). All of the observed complications in the present study were related primarily to wound break-

down or discomfort associated with the subcutaneous penetration of the extension cable. The dermal and subcutaneous layers in this area are generally thin, and this undoubtedly contributed to the observed complication rate. The recent availability of smaller connectors with a lower profile makes it likely that this complication rate will decrease in the near future.

## CONCLUSIONS

These findings indicate that peripheral stimulation of the supraorbital or infraorbital branches of the trigeminal nerve may be an effective method for relief of neuropathic facial pain occurring after trauma or herpes zoster infection in a majority of patients. A suggested method of this novel approach to the treatment of these disorders is thus warranted.

## REFERENCES

1. Appleby DC, Lewis PH: Treatment of postherpetic neuralgia: A systematic review of the literature. *Ann Intern Med* 127:117-124, 1997
2. Johnson J: Herpetic trigeminal neuropathic pain: A review. *Acta Neurol Scand Suppl* 17:23-37, 1999
3. Jannat K: A new classification of facial pain. *Diagnosis and Management in Neurosurgery* 5:1: 30, 2002
4. Johnson J, Anderson M, Brown H, Fowler B, Greenman S, Gentry G, Gofner R, Hickey J, Johnson D, Prosser W, et al: Herpes zoster ophthalmicus: clinical and genetic risk factors and genetic susceptibility. *Proc Natl Acad Sci USA* 93:2754-2758, 1996
5. Dworkin R: A comparison of neuropathic pain, Spondylosis, neuropathic, and central mechanisms. *Clin J Pain* 14:944-950, 2002
6. Elia S, Kim D, Grewal V, Brown D, Tamura S: Central processing mechanisms in trigeminal neuropathic pain. *Acta Neurol Scand* 101:95-100, 2000
7. Kato H, Nakase T: Trigeminal neuropathic pain: Pathophysiology and mechanisms: A quantitative assessment of abnormal pain and sensory neuropathy. *Neurology* 47:1144-1150, 1997
8. McMahon SB
9. Hirsch J, Ferreira M: Postherpetic neuralgia: surgery and facial neurectomy. *Neural Res* 12:117, 1997
10. Kasperk J, Johnson R, Grewal M: Treatment of postherpetic neuropathic pain. *Comp Ther* 20:115-119, 1996
11. Kopylov V, Grewal V, Akay S, Taylor S: CT-guided, 30% procaine neurolytic blockade. *Neurosurgery* 36:94-98, 1994
12. Lane JD, Swain J, Kishitani D: Electrocutaneous analysis of 22 patients with chronic pain treated by peripheral nerve stimulation. *J Neurosurg* 73:452-458, 1991
13. Newman BA, Lundberg M, Johnson R, Jannat K: Postherpetic trigeminal neuropathic pain: treatment of trigeminal neuropathic pain. *Acta Neurol Scand* 101:101-107, 2000
14. Pappagallo G, Grolander AL, Suthana P, Vanni J, de Marco MR, Raju JN: Heterogeneous pattern of sensory evoked potentials in postherpetic neuralgia suggest multiple pathophysiological mechanisms. *Anesthesiology* 102:911-924, 2005
15. Pappagallo G, Schwartz J, Suthana P, Grolander AL, de Marco MR, Raju JN: Heterogeneous pattern of sensory evoked potentials in postherpetic neuralgia suggest multiple pathophysiological mechanisms. *Anesthesiology* 102:911-924, 2005
16. Johnson J
17. Johnson J, Brown M, Grewal R: Chronic postherpetic neuralgia of the gasserian ganglion for the relief of pain in a series of 14 patients. *J Neurosurg* 1: 949-957, 1997
18. Tsubokawa T, Nakayama Y, Yamamoto T, Horiuchi T, Koyama S: Clinical efficacy of laser stimulation for the treatment of central pain. *Acta Neurol Scand Suppl* 161:10-14, 1993
19. Under-States M, Cameron S, Lee D: The use of stimulation of postherpetic neuralgia. *Can J Neuro Sci* 1:129-135, 2001

*E. Vinken, J.C. Haenen, M.P. Neumeister and J. van't Hof-Jensen are pathophysiology Dept Editor, J 24 25, 26, 27, 28*

## COMMENTS

It is well known that painful trigeminal neuropathy is extremely difficult to manage, and most of the treatment modalities available rarely are of much assistance. These modalities are not uncommon and may occur as result of frequently performed facial surgery, for example, of the maxillary sinus (Marswell's procedure). Local injections of alcohol or, as demonstrated, even a potentially effective therapy for this condition, direct stimulation of the gasserian ganglion and infraorbital trigeminal nodules often was attempted. In some patients, such stimulation could provide effective pain relief for many years, but it failed in the majority of patients. Furthermore, the placement of the stimulation electrode with a subdermal approach required major surgery, and, because of the location of the point, possibly implanted electrode in the gasserian system often was problematic. Therefore, the novel stimulation approach presented in this article is of much interest. Although the number of patients is small, the follow-up period is relatively long, and the outcome is positively favorable. A possible drawback is that this method, at least at this stage, seems to be limited to first- and second-branch neuralgia.

It is somewhat surprising that the concept of peripheral nerve stimulation has not been applied previously to the trigeminal nerve. In the early literature regarding transcutaneous electrical nerve stimulation, there are reports that transcutaneous stimulation was applied for various forms of facial pain. In the late 1970s, when we developed the technique for direct trigeminal stimulation, the patients were screened with transcutaneous electrical nerve stimulation. In addition to the discomfort of having a large rubber-plate taped to their face, however, several of the patients experienced aggravated cutaneous allodynia in the form of allodynia, hyperalgesia, and so forth as a result of the stimulation.

In the present study, pain relief is assessed as a percentage in a global evaluation. Trigeminal neuropathy usually is associated with profound changes of cutaneous sensitivity, which gives rise to evoked pain in addition to ongoing, spontaneous pain components. Both gasserian stimulation and motor cortex stimulation effectively suppressed the cutaneous signs of neuropathy, and it would have been interesting if the same effects were observed with peripheral trigeminal stimulation. This seems to be the case. A detailed assessment with quantitative sensory testing would have permitted analysis of and insight in possible mechanisms of action of the stimulation.

It is not surprising that the outcome tended to be less positive in patients presenting with postherpetic neuralgia than in those with a history of trigeminal trauma, considering the complexity of the latter condition. There is evidence that the form and extent of surgically changes in postherpetic neuralgia might be different pain-generating mechanisms, hence, presumably also to the likelihood of response to the

stimulation. It might therefore be that a thorough analysis of these changes could help to predict the responsiveness. The probability of alleviating pain associated with trigeminal neuropathy by use of peripheral stimulation, and the simplicity of the implantation procedure as compared with motor cortex stimulation, make this an attractive treatment option.

Björn A. Meyerson  
Stockholm, Sweden

The authors provide their extensive data for 11 patients with facial neuropathic pain, including trigeminal postherpetic neuralgia, trigeminal posttraumatic neuropathic pain, and bilateral trigeminal neuralgia, who underwent trigeminal occipital stimulation. Satisfactory pain relief was obtained in 73% of patients. Although peripheral stimulation has been used previously in the treatment of occipital neuralgia, the reported efficacy of trigeminal peripheral stimulation in patients with facial pain and the technical aspects of the procedure are original. This study demonstrates promising results of this new treatment modality in patients with intractable facial pain, whose management presents great difficulty. Stimulation of the trigeminal nerve is not a new concept, however, and gasserian and subgasserian neurostimulation have been performed by several authors since 1960 (1, 2). Although the technique described is a non-peripheral approach, these procedures may be considered as a group.

The efficacy of such procedures is not well understood. Neuropathic pain is known to be a result of pathological changes in the central nervous system. Surgical procedures usually are performed on targets within the central nervous system such as the dorsal root entry zone, substantia gelatinosa, trigeminal tract, nucleus, and motor cortex. Peripheral interventions principally are avoided, as they may increase central neuronal degeneration. The beneficial effect of very peripheral stimulation of the trigeminal nerve is obvious but surprising. The selection criteria for stimulation of the cranial part of the trigeminal nerve are unclear, and a rationale is needed for choosing peripheral rather than gasserian or subgasserian stimulation. Finally, technical difficulty in stimulating the third branch of the trigeminal nerve seems to be a disadvantage of trigeminal peripheral stimulation.

Transcutaneous trigeminal tractotomy-neurotomy also should be considered in patients with facial neuropathic pain of various causes, including postherpetic neuralgia (2). We performed this procedure in 53 cases, and complete or satisfactory pain control was obtained in 77.0% of cases. Four of patients had postherpetic neuralgia, and complete pain was obtained in two patients. In a third patient, pain relief was unsatisfactory, a trigeminal dorsal root entry zone operation was performed, and complete pain relief was obtained. Trigeminal tractotomy-neurotomy was not effective in the fourth patient. This procedure is efficient and minimally invasive. Although it has been classified as a neurologically destructive procedure, it is associated with a low rate of complications. Conversely, neurostimulation is a technologically

sophisticated treatment methods, if they increase the patient's dependence on the physician and hospital, and if it is expensive.

Trigeminal peripheral stimulation is a new surgical treatment method in patients with focal neuropathic pain. The authors present an interesting, original, and efficient treatment modality. However, the mechanism of its efficacy needs to be explained with physiologic studies, and selection criteria of the trigeminal nerve neurectomy event. Long-term follow-up should be a mandatory established with further studies. Larger series with longer follow-up periods are needed to validate the decision to conduct trigeminal nerve stimulation in patients with intractable facial pain.

Yücel Kaşpıralı  
Ali Savas  
Ankara, Turkey

1. Elstermann G, Vennart H. Carotid Reafferentation (glucose injection in percutaneous carotid artery) in the treatment of typical and atypical trigeminal pain. *Neural Reg* 1994; 14: 1987.
2. Kınacı M, Savaş A, Çaylıoğlu B, Ayvan M, Taşoğlu M, Ayvan E. Cefalodin percutaneous intracavernous neurectomy. *Türk Neurology* 5 Oct 2011; 197A.
3. Elstermann G, Aronowart H. The effects of intracavernous stimulation of the carotid ganglion for treatment of typical facial neuropathic. *Phong Clin Neurophysiol* 2007; 118: 1897.

**T**reatment of facial pain caused by trigeminal neuralgia has an extremely good neurosurgical track record. The results of surgery for trigeminal postherpetic neuropathy and posther-

petic pain, however, are dismal. That treatment for neuropathic pain in the territory of the trigeminal nerve should have such disparate results depending on the origin of the neurostatic process is vexing for any neurosurgeon who treats such patients. The lack of neurosurgical options for patients with trigeminal neuropathy and postherpetic pain seems to be changing with the publication of reports such as the present study. The authors describe a novel application of peripheral nerve stimulation to isolated V<sub>1</sub> or V<sub>2</sub> trigeminal pain caused by postherpetic neuralgia or nerve injury. The surgical procedures described in this manuscript are straightforward and are associated with low morbidity. The quality of the data collection in this retrospective study seems high, given the usual caveats of retrospective studies. There was an 80% direct data acquisition rate, with the remaining 20% of data gleaned from charts. This excellent data collection rate for this type of study rather guarantees the quality of the data. The follow-up period is relatively long, although a low number of patients (which is fairly small). A larger retrospective series would allow clinicians to determine a better idea the precise symptoms best alleviated by use of these devices. Prospective study of this surgical intervention should prove fairly straightforward given the intractable nature of these disease processes and clear clinical criteria to define them.

Oren Sapher  
Ann Arbor, MI

#### Future Meetings—Congress of Neurological Surgeons

The following are the planned sites and dates for future annual meetings of the Congress of Neurological Surgeons:

|      |                   |                 |
|------|-------------------|-----------------|
| 2004 | San Francisco, CA | October 14–21   |
| 2005 | Aspen, CO         | October 8–13    |
| 2006 | Chicago, IL       | October 24–27   |
| 2007 | San Diego, CA     | September 14–21 |

#### Future Meetings—American Association of Neurological Surgeons

The following are the planned sites and dates for future annual meetings of the American Association of Neurological Surgeons:

|      |                   |                  |
|------|-------------------|------------------|
| 2005 | New Orleans, LA   | April 16–21      |
| 2006 | San Francisco, CA | April 22–27      |
| 2007 | Washington, DC    | April 13–19      |
| 2008 | Chicago, IL       | March 26–April 1 |